

MONITORING HEAVY METALS IN THREE REGIONS OF ITALY USING MOSSES AND SOILS AS INDICATORS: A MULTIVARIATE AND GIS APPROACH



¹Gramatica, P., ¹Giani, E., ¹Papa, E., ²Preatoni, D., ³Jones, R. J. A., ⁴Musmeci, L. and ³Cenci, R. M.

¹QSAR and Environmental Chemistry Research Unit, University of Insubria, DBSF, Varese (Italy) - ²Biocoenosis Analysis and Management Unit, University of Insubria, DASS, Varese (Italy)- Institute for Environment and Sustainability - ³Joint Research Centre, Ispra (Italy) - ⁴Italian National Institute of Health, Rome, Italy

Web: <http://dipbsf.uninsubria.it/qsar/> e-mail: paola.gramatica@uninsubria.it

ABSTRACT

A biomonitoring survey involving the moss *Hypnum cupressiforme* Hedw. was carried out in three Italian regions, Piedmont, Aosta Valley and Sicily, to evaluate the atmospheric deposition of the elements Cd, Cu, Hg, Ni, Pb and Zn. Soil samples were taken from the same areas and these were analysed for the same elements. Heavy metal concentrations in mosses have permitted the evaluation of the deposition rates of the studied elements. These concentrations were found to be significantly higher in Piedmont than in Sicily because of the higher level of industrialisation in Piedmont. The combined approach using soils and mosses led to the discrimination, by the enrichment factor (on Al basis), of the anthropogenic and pedological/geological sources. The fallout of elements Hg and Zn in Piedmont, Hg in Aosta Valley, Cd and Hg in Sicily were found to be predominantly anthropogenic in origin, whereas other metals are predominantly of pedological/geological origin. Moss and soil concentration data were analysed using Principal Components Analysis (PCA). This multivariate analysis permitted a visual representation of the co-distribution of metals in the mosses and soil, and an evaluation of the site similarities in metal composition. The results are also presented in map form using the kriging approach integrated in the Geographic Information System (GIS).

INTRODUCTION

The emergency created by environmental pollution, caused by human activities (both industrial and domestic), means that tools are needed to determine the quality and quantity of pollutants in the environment. Recording systems are necessary to identify appropriate strategies of reduction of pollutants emitted and to verify, afterwards, the effectiveness or the inadequacy of measures taken. Technological research has provided and improved instrument systems to record air, soil, water and biota pollution. Since these instruments are often characterized by high cost just few of them are usually applied; moreover being these techniques projected to record only certain substances, they are unable to detect the presence of new pollutants. In order to complete the information given by the technical instruments there is a spreading use of living organisms to detect the presence of pollutants and to determine their effect on live communities.

MATERIALS & METHODS

Data: Soil and mosses samples were collected in 28 sites in Sicily, in 41 sites in Piedmont and in 20 sites in Aosta Valley.

The methodology adopted for the samples analysis was applied according to the A.N.P.A. method [1]. The heavy metals Cd, Cu, Hg, Ni, Pb and Zn were investigated.

Principal Components Analysis (PCA): linear combinations of the input variables (metal concentration in this case) to new uncorrelated variables, called principal components, which account for the variance in the data. The first PC explains the main part of the variance of the original data set.


Geostatistical Methods and Mapping: each data set was transformed in logarithmic units. The heavy metal levels were shown on contour maps using the coordinates of sampling sites. Maps were obtained using an universal kriging interpolator [2].

RESULTS


Principal Components Analysis (PCA) of metal concentrations in Aosta Valley and Piedmont mosses and soils was performed in order to have a multivariate view and a graphic representation of the overall metal distribution in mosses and soils in the studied sites (Fig.1-2) identifying the most polluted zones. The normalization of the concentration values of heavy metals in soils and mosses, considering Al as the conservative element, allows the discrimination between anthropogenic and natural heavy metals' sources using the following formula: Enrichment Factor (EF) = [(X moss/Al moss)/(X soil/Al soil)]

EF ≥ 10: heavy metal concentrations observed in moss could be originated by anthropogenic activity and/or natural events such as volcanic eruptions and large forest fires.

EF < 10: heavy metal concentrations observed in moss are related to their concentration in soil and/or substrate (natural sources);



The PCA of metal concentrations in Sicily, Aosta Valley and Piedmont mosses and soils (Figures 1-2) show that the Aosta Valley and Piedmont sites have higher metal concentrations than Sicily sites, except for Pollina in fig.1 and for Vizzini and Salemi in fig. 2. The EF analysis of the mosses and soils for the same site allows the identification of the sites polluted by specific metals of anthropogenic origin.



The GIS approach was adopted to distinguish the sites characterized by high concentration of Cd, Cu, Hg, Ni, Pb and Zn in the three Regions. Some results are presented in the form of coloured maps reported.

CONCLUSIONS

> Mosses are good indicators of atmospheric pollution by heavy metals useful for the monitoring at a regional scale, for the estimation of the actual contamination or of the temporal changes in contamination.

> PCA performed on heavy metals concentrations measured in Sicily, Piedmont and Aosta Valley sites, identified the most polluted sites and the similarity in pollution typology.

> GIS mapping is useful for immediate visualization of metal distribution and contamination level in a region.




Figure 3 - Cd, Hg, Pb and Zn in soils

Figure 4 - Ni, Pb and Zn in mosses

REFERENCES

- [1] R.M. Cenci. 1999. Agenzia Nazionale per la Protezione dell'Ambiente. Biomonitoraggio della qualità dell'aria sul territorio nazionale. Serie Atti 2/1999. 241-263;
- [2] P.A. Burrough e R.A. McDonnell. (1998). Oxford. pp. 333.