

Robert SUPPER and Ivo BAROŇ (Eds.)

Landslide Monitoring Technologies & Early Warning Systems

Current Research and Perspectives for the Future



Book of extended abstracts
Open Workshop within the frame of the EU FP7 "SafeLand" Project
February 24th, 2010, Vienna

Berichte der Geologischen Bundesanstalt, 82

Landslide Monitoring Technologies & Early Warning Systems

Current Research and Perspectives for the Future

Book of extended abstracts
Open Workshop within the frame of the EU FP7 "SafeLand" Project
February 24th, 2010, Vienna

Editors: Robert SUPPER and Ivo BAROŇ

Berichte der Geologischen Bundesanstalt, Nr. 82

Wien, August 2010

Table of Content

Introductory Foreword	7
The SafeLand Consortium, N. CASAGLI & R. SUPPER: The Safeland Project – General Overview and Monitoring Technology Development	9
BAROŇ, I. & SUPPER, R.: State-of-the-Art of Landslide Monitoring in Europe: Preliminary Results of the SafeLand Questionnaire	19
SCHIANO, P: Meteorological and Climate Forecasting for Landslide Prediction	26
MICHOUD, C., DERRON, M.-H., ABELLÁN, A. & JABOYEDOFF, M.: A Review of Remote Sensing and Ground-Based Techniques for Landslides Detection, Fast Characterization, Rapid Mapping and Long-Term Monitoring	27
TRAVELLETTI, J., MALET, J.P., SCHMITTBUHL, J., TOUSSAINT, R., DELACOURT, C. & STUMPF, A.: A Multi-Temporal Image Correlation Method to Characterize Landslide Displacements	29
KERLE, N., STUMPF, A. & MALET, J-P.: Object-Oriented and Cognitive Methods for Multi-Data Event-Based Landslide Detection and Monitoring	33
TOFANI, V., CATANI, F. & CASAGLI, N.: Weather Forecasting and Radar Technologies for Landslide Prediction and Mapping: Some Examples in Italy	35
VAN DEN EECKHAUT, M. & HERVÁS, J.: Testing Different Techniques for Detection, Rapid Mapping and Monitoring of Landslides in the Barcelonnette Region Using Satellite and Airborne Optical Imagery	38
PICARELLI, L., FORNARO, G. & COMEGNA, L.: Advanced Criteria and Techniques for Landslide Monitoring	40
CASCINI, L., PEDUTO, D. & FORNARO, G.: A New Approach to the Use of DInSAR Data in Landslide Studies at Different Scales: Case study of National Basin Authority of Liri-Garigliano and Volturno rivers (Italy)	43
ČARMAN, M. & KUMELJ, S.: GeoZS - Landslide Issues and Contributions to the SafeLand – The Stovze Landslide	45
GILI, J.A.: DInSAR vs. Wire Extensometer Calibration, and GBSAR First Survey at the Vallcebre Landslide (Spain)	47
AGLIARDI, F., CROSTA, G.B. & FRATINI, P.: Forecasting the Failure of Large Landslides for Early Warning: Issues and Directions	50
SUPPER, R., BAROŇ, I., JOCHUM, B., ITA, A., MOTSCHKA, K. & WINKLER, E.: Airborne Geophysics and Geoelectric and Inclinoetric Monitoring at the Gschlifgraben Landslide	52
LOVISOLO, M.: Bagnaschino Landslide: from Early Warning to Site-Specific Kinematic Analysis	59
CARDELLINI, S. & OSIMANI, P.: Living with Landslides: the Ancona Case History and Early Warning System	63
ROTH, M. & BLIKRA, L.H.: Seismic Monitoring of the Unstable Rock Slope at Åknes, Norway	69

Testing Different Techniques for Detection, Rapid Mapping and Monitoring of Landslides in the Barcelonnette Region Using Satellite and Airborne Optical Imagery

M. VAN DEN EECKHAUT & J. HERVÁS

Land Management and Natural Hazards Unit, Institute for Environment and Sustainability Joint Research Centre (JRC) – European Commission, 21027 Ispra (VA), Italy, miet.van-den-eeckhaut@jrc.ec.europa.eu, javier.hervas@jrc.ec.europa.eu

Within SafeLand the Joint Research Centre (JRC) of the European Commission is involved in three research areas, i.e. Area 2 on "Quantitative risk assessment", Area 3 on "Quantifying global change scenarios (climatic and anthropogenic) and their impact on land-slide hazard and risk in the future" and Area 4 on "Development of monitoring technology, especially early warning systems and remote sensing techniques, and applications". This abstract focuses on the activities within Area 4. A short overview will be provided of the research planned by JRC. For more elaborate information on the techniques presented, reference is made to some clarifying research articles.

With the increasing spatial resolution of e.g. IKONOS (1999), QuickBird (2001), OrbView-3 (2003), now phased out, Worldview (2007; 0.44 m pan) or GeoEye (2008; 0.41 m pan) sensors, application fields which had previously been the domain of airborne remote sensing (ARS) could be tackled by satellite remote sensing (SRS [BLASCHKE, 2010]). Indeed, two of the three methods that will be tested and eventually improved by JRC within Area 4 have been originally designed with ARS images. JRC will test two techniques for detection and rapid mapping of landslides and one technique for long-term change detection of landslides. The techniques are planned to be applied to very high resolution (VHR) spaceborne optical sensors and airborne LiDAR. For this, the Barcelonnette test site (200 km²; Alpes-de-Haute-Provence, France) was selected, but probably also data from other SafeLand test sites will be used.

With regard to landslide detection and mapping, the first technique to be tested and eventually improved is a semi-automatic texture classification technique developed by HERVÁS & ROSIN (1996), HERVÁS et al. (1996) and HERVÁS & ROSIN (2001). These authors obtained good results with a discrimination method for landslide mapping in a semi-arid, sedimentary area in SE Spain from high resolution Daedalus ATM (3 m), IRS-1C Pan (5 m) and SPOT Pan (10 m) images. The second landslide identification technique will use LiDAR data in combination with optical images for Object Oriented Analysis (OOA; e.g. MARTHA et al., 2010). The possibilities of the Definiens Developer software for segmentation and classification of landslides are illustrated with results obtained from the exploration of the software.

For long-term detection of landslide activity, a method entailing automatic change detection of suitably pre-processed multitemporal images, followed by thresholding of pixel intensities into landslide-related change pixels was developed by HERVÁS et al. (2003). The method was previously applied to VHR (1 m) orthoimages of the Tessina landslide (Italy) and provided representative results on the pixel intensity changes related to the reactivation of the landslide in 1992 (Figure 1).

In agreement with SafeLand's objective to secure data exchange with other Areas, the results obtained from this study will be used as input data in Areas 2 and 3.




Fig. 1: Surface changes in the Tessina landslide between 1988 and 1994, illustrated on the 1994 orthophotograph. (1) within the landslide body, positive pixel intensity changes represent new soil outcrops and remobilised soil as a result of landslide reactivation; (2) negative pixel intensity changes are due to vegetation growth or soil moisture increase. Most changes outside the landslide correspond to land use change. Coordinates in UTM (HERVÁS et al., 2003).

References

- BLASCHKE, T. (2010): Object based image analysis for remote sensing. *ISPRS Journal of Photogrammetry and Remote Sensing*, 65, 2–16.
- HERVÁS, J. & ROSIN, P.L. (1996): Landslide mapping by textural analysis of Daedalus ATM data. *Proc. 11th Thematic Conf. Applied Geologic Remote Sensing*, Las Vegas, Nevada, 2, ERIM, Ann Arbor, Michigan, USA, 394–402.
- HERVÁS, J. & ROSIN, P.L. (2001): Tratamiento digital de imágenes de teledetección en el espectro óptico para el reconocimiento y control de deslizamientos. *Proc. V Simposio Nacional sobre Taludes y Laderas Inestables*. Madrid, 27–30 November 2001. CEDEX, Ministerio de Fomento, Madrid, 63–74.
- HERVÁS, J., ROSIN, P.L., FERNÁNDEZ-RENAU, A., GÓMEZ, J.A. & LEÓN, C. (1996): Use of airborne multispectral imagery for mapping landslides in Los Vélez district, south-eastern Spain. In: CHACÓN, J., IRIGARAY, C. & FERNÁNDEZ, T. (Eds.), *Landslides*. Balkema, Rotterdam, 353–361.
- HERVÁS, J., BARREDO, J.I., ROSIN, P.L., PASUTO, A., MANTOVANI, F. & SILVANO, S. (2003): Monitoring landslides from optical remotely sensed imagery: the case history of Tessina landslide, Italy. *Geomorphology*, 54, 63–75.
- MARTHA, T.R., KERLE, N., JETTEN, V., VAN WESTEN, C. & KUMA, K.V. (2010): Characterising spectral, spatial and morphometric properties of landslides for semi-automatic detection using object-oriented methods. *Geomorphology* 116, 24–36.