SOILS OF THE GREAT PLAINS

SOIL LEGEND

UDIC BOROLLS AND AQUOLLS

1. Argiborolls-Eutroboralfs; undulating-rolling; fine-loamy and clayey
2. Argiborolls-Haplaquolls; level-undulating; clayey, fine-loamy and fine-silty
3. Argiborolls-Haploborolls; level-undulating; fine-loamy
4. Argiborolls-Haploborolls; undulating-hilly; fine-loamy
5. Argiborolls-Haploborolls-Natriborolls; level; clayey and fine-silty
6. Calciaquolls; level; fine-silty; saline
7. Calciaquolls-Haploborolls; level; coarse-silty and fine-silty
8. Calciaquolls-Haploborolls-Argialbolls; level; fine-loamy and clayey
9. Haplaquolls-Calciaquolls; level; clayey and fine-silty; vertic
10. Haploborolls; level; fine-loamy over sandy or sandy-skeletal and fine-loamy
11. Haploborolls; level-undulating; fine-silty
12. Haploborolls; undulating-rolling; fine-loamy
13. Haploborolls-Argiborolls; level-undulating; fine-silty and fine-loamy
14. Haploborolls-Calciaquolls; level-undulating; coarse-loamy
15. Haploborolls-Calciaquolls; level-undulating; fine-loamy
16. Haploborolls-Calciaquolls-Haplaquolls; level; coarse-loamy and sandy
17. Haploborolls-Haplaquolls; level-undulating; fine-silty and fine-loamy
18. Natriborolls; level-undulating; clayey and fine-loamy

TYPIC BOROLLS AND USTROTHENTS

19. Argiborolls; level-undulating; fine-loamy
20. Argiborolls-Argialbolls-Haploborolls; level-undulating; fine-loamy and clayey
21. Argiborolls-Haploborolls; level-rolling; fine-silty and fine-loamy
22. Argiborolls-Haploborolls; level-hilly; clayey and clayey-skeletal; lithic
23. Argiborolls-Haploborolls-Ustorthents; level-rolling; fine-loamy
24. Argiborolls-Natriborolls-Ustorthents; level-rolling; loamy and clayey
25. Argiborolls-Ustorthents; level-rolling; loamy
26. Argiborolls-Ustorthents; level-rolling; fine-loamy
27. Argiborolls-Ustorthents; level-rolling; clayey and fine-loamy
28. Calciborolls-Argiborolls; undulating-rolling; fine-loamy, clayey and loamy-skeletal
29. Calciborolls-Calciorthids; level-rolling; fine-loamy and loamy-skeletal
30. Haploborolls-Argiborolls-Ustipsamments; level-rolling; loamy and sandy
31. Haploborolls-Ustorthents-Argiborolls; undulating-hilly; fine-loamy and loamy-skeletal; lithic
32. Haploborolls-Ustorthents-Argiborolls; undulating-hilly; fine-loamy
33. Haploborolls-Ustorthents-Ustochrepts; undulating-hilly; fine-silty and fine-loamy
34. Ustorthents-Argiborolls; undulating-hilly; fine-loamy
35. Ustorthents-Argiborolls; undulating-steep; loamy; shallow

ARIDIC BOROLLS AND TORRIORTHENTS

36. Argiborolls; level-undulating; clayey
37. Argiborolls-Haploborolls-Paleargids; level-undulating; clayey and loamy
38. Argiborolls-Paleargids-Natrargids; level-undulating; clayey
39. Argiborolls-Torriorthents; undulating-rolling; clayey and fine-loamy
40. Haploborolls-Eutroboralfs-Argiaborolls; undulating-hilly; coarse-loamy and fine-loamy
41. Torriorthents; undulating-steep; clayey, vertic; shallow
42. Torriorthents-Argiborolls; level-undulating; clayey; vertic

BOROLLIC ARIDISOLS AND TORRIORTHENTS

43. Natrargids; level; clayey
44. Torriorthents-Camborthids-Natrargids; undulating-hilly; loamy and clayey; shallow

UDOLLS, MESIC

45. Argiudolls; level-rolling; clayey and fine-loamy
46. Argiudolls-Argiaquolls-Argialbolls; level; clayey
47. Argiudolls-Hapludolls; level-rolling; clayey, fine-silty and fine-loamy
48. Argiudolls-Hapludolls; level-rolling; clayey and fine-silty
49. Hapludolls-Udorthents-Hapludalfs; undulating-hilly; fine-silty

UDIC USTOLLS AND AQUOLLS, MESIC

50. Argiustolls; level-rolling; clayey and fine-silty
51. Argiustolls-Argiaquolls-Argialbolls; level-undulating; clayey
52. Argiustolls-Haplustolls; level-rolling; clayey and fine-silty
53. Argiustolls-Haplustolls; level-steep; clayey-skeletal, clayey and loamy; lithic
54. Argiustolls-Haplustolls, lithic, loamy; undulating-hilly; clayey, fine-loamy and fine-silty
55. Argiustolls-Natrustolls; level-undulating; clayey
56. Argiustolls-Pellusterts-Haplustolls; level-undulating; clayey
57. Haplaquolls-Argiustolls-Ustifluvents; level; clayey and loamy
58. Haplustolls; level-undulating; fine-silty and fine-loamy
59. Haplustolls; level-rolling; fine-silty
60. Haplustolls-Ustipsamments; level-rolling; coarse-silty, coarse-loamy and sandy
61. Haplustolls-Ustipsamments; undulating-rolling; sandy and coarse-loamy
62. Haplustolls-Ustorthents; level-rolling; fine-silty
63. Haplustolls-Ustorthents; level-hilly; loamy and sandy
64. Haplustolls-Ustorthents; undulating-hilly; fine-silty

TYPIC USTOLLS AND AQUOLLS, MESIC

65. Argiustolls; level-undulating; clayey
66. Argiustolls-Argiaquolls; level-undulating; fine-silty and clayey
67. Argiustolls-Haplustolls; level-undulating; fine-loamy and fine-loamy over sandy or sandy-skeletal
68. Argiustolls-Haplustolls; level-rolling; fine-silty and fine-loamy
69. Argiustolls-Haplustolls; level-rolling; clayey; vertic
70. Argiustolls-Haplustolls-Calciustolls; level-rolling; clayey, fine-silty and fine-loamy
71. Argiustolls-Haplustolls-Natrustolls; level-rolling; fine-loamy and clayey
72. Argiustolls-Haplustolls-Ustorthents; level-rolling; fine-silty
73. Argiustolls-Natrustolls; level-undulating; clayey and fine-loamy
74. Argiustolls-Natrustolls; level-rolling; clayey and fine-silty
75. Calciustolls-Argiustolls-Haplustolls; undulating-hilly; loamy and clayey; shallow
76. Haplaquolls-Haplustolls-Ustipsamments; level-undulating; sandy
77. Haplustolls; level-rolling; clayey; vertic
78. Haplustolls-Argiustolls-Ustorthents; undulating-rolling; fine-loamy
79. Haplustolls-Argiustolls-Ustorthents; shallow, loamy; level-hilly; coarse-loamy, sandy, fine-loamy over sandy or sandy-skeletal
80. Haplustolls, shallow-Argiustolls; undulating-steep; clayey and loamy
81. Haplustolls-Ustipsamments; undulating-rolling; sandy, coarse-loamy, and fine-loamy over sandy or sandy-skeletal
82. Haplustolls-Ustorthents; level-hilly; loamy and sandy
83. Haplustolls-Ustorthents; undulating-hilly; fine-loamy
84. Haplustolls-Ustorthents, shallow; undulating-steep; clayey; vertic
85. Haplustolls-Ustorthents-Argiustolls; undulating-hilly; fine-silty

ARIDIC USTOLLS, USTALFS, AND TORRIORTHENTS, MESIC

86. Argiustolls; level-undulating; fine-loamy and fine-loamy over sandy or sandy-skeletal
87. Argiustolls-Haplargids Paleustolls; level-rolling; fine-loamy, clayey and coarse-loamy
88. Argiustolls-Haplustolls; level-undulating; fine-silty and clayey
89. Argiustolls-Haplustolls; level-rolling; clayey and loamy; lithic
90. Argiustolls-Haplustolls; undulating-rolling; coarse-loamy, fine-loamy over sandy or sandy-skeletal, sandy and fine-loamy
91. Argiustolls-Haplustolls-Haploborolls; undulating-hilly; loamy-skeletal and loamy
92. Argiustolls-Haplustolls-Torriorthents; level-undulating; clayey and fine-silty
93. Argiustolls-Haplustolls-Torriorthents; level-rolling; fine-silty
94. Argiustolls-Haplustolls-Torriorthents; undulating-rolling; loamy
95. Argiustolls-Haplustolls-Torripsamments; undulating-rolling; fine-loamy, coarse-loamy and sandy
96. Argiustolls-Natrustalfs-Pellusterts; level-undulating; coarse-loamy and clayey
97. Argiustolls-Torriorthents; undulating-hilly; fine-silty
98. Argiustolls-Torriorthents, lithic, loamy; undulating-rolling; fine-loamy and clayey
99. Argiustolls-Torriorthents, shallow-Camborthids, undulating-hilly; clayey and loamy
100. Haplustolls-Paleorthids; level-rolling; fine-loamy; lithic
101. Haplustolls-Torriorthents; level-rolling; loamy
102. Haplustolls-Torriorthents; undulating-hilly; fine-silty
103. Haplustolls-Torriorthents, shallow, loamy-Torripsamments; undulating-hilly; coarse-loamy, coarse-silty, and sandy
104. Paleustalfs-Haplustalfs-Ustipsamments; undulating-rolling; fine-loamy, coarse-loamy, and sandy
105. Paleustolls-Argiustolls-Paleorthids; level-undulating; clayey and loamy; lithic
106. Paleustolls-Calciustolls; level-undulating; clayey and fine-loamy
107. Paleustolls-Haplargids-Torriorthents; level-rolling; clayey, fine-loamy, and fine-silty

USTIC ARIDISOLS AND TORRIORTHENTS, MESIC

108. Calciorthids-Torriorthents; level-undulating; fine-loamy and loamy-skeletal; saline
109. Camborthids-Argiustolls; undulating-rolling; clayey and loamy; vertic
110. Camborthids-Torriorthents; level-undulating; clayey
111. Camborthids-Torriorthents, shallow; undulating-hilly; clayey; vertic
112. Haplargids-Calciorthids; level-rolling; fine-silty, fine-loamy, and coarse-loamy
113. Haplargids-Calciorthids-Haplustolls; level-undulating; fine-silty, and fine-loamy
114. Haplargids, fine-loamy-Calciorthids-Paleorthids; level-undulating; loamy; shallow
115. Haplargids-Torrifluvents, loamy; level-undulating; fine-silty and clayey
116. Haplargids-Torriorthents; level-undulating; clayey and fine-silty
117. Haplargids-Torriorthents-Natrargids; undulating-rolling; clayey and loamy
118. Haplargids-Torripsamments; level-undulating; coarse-loamy, fine-loamy, and sandy
119. Paleorthids-Calciorthids-Haplargids, fine-loamy; level-rolling; loamy; shallow
120. Torriorthents; undulating-hilly; loamy; lithic
121. Torriorthents; rolling-steep; loamy and clayey; shallow
122. Torriorthents, lithic, loamy-Haplargids-Argiustolls; undulating-hilly; clayey and fine-silty
123. Torriorthents-Haplargids-Natrargids; undulating-hilly; loamy and clayey; shallow

UDOLLS, THERMIC

124. Argiudolls-Argiaquolls-Haplustolls, lithic, loamy; level-hilly; clayey and fine-silty
125. Argiudolls-Haplustolls, lithic, loamy-Paleudolls; undulating-hilly; fine-silty, clayey, clayey-skeletal
126. Paleudolls-Albaqualfs-Argiudolls; level-rolling; clayey and fine-loamy

UDIC USTOLLS, USTALFS, AND AQUALFS, THERMIC

127. Albaqualfs-Paleustalfs; level-undulating; clayey
128. Haplustolls-Ustochrepts; undulating-hilly; clayey and loamy; lithic
129. Haplustolls-Ustochrepts; undulating-hilly; clayey-skeletal, loamy, and clayey; lithic
130. Ochraqualfs-Paleustalfs-Pellusterts; level-undulating; clayey
131. Paleustalfs-Albaqualfs-Pellusterts; level-undulating; clayey and loamy
132. Paleustalfs-Haplustalfs-Ustochrepts, shallow; undulating-rolling; clayey, loamy, and sandy
133. Paleustolls-Argiustolls-Ustochrepts; level-rolling; clayey, fine-silty, and fine-loamy

TYPIC USTOLLS, USTALFS, AND USTOCHREPTS, THERMIC

134. Argiustolls-Argiaquolls, clayey; level-undulating; fine-loamy and coarse-loamy
135. Argiustolls-Calciustolls-Paleustalfs; level-undulating; clayey and fine-loamy
136. Argiustolls-Haplustolls; level-undulating; fine-silty and coarse-silty
137. Argiustolls-Paleustolls-Ustochrepts; level-undulating; clayey
138. Calciustolls-Paleustolls; level-undulating; clayey; lithic
139. Haplustalfs-Argiaquolls, clayey-Ustipsamments; level-undulating; sandy and coarse-loamy
140. Haplustalfs-Ustipsamments-Ustorthents; level-rolling; sandy and coarse-loamy
141. Haplustolls-Calciustolls-Argiustolls; level-hilly; clayey-skeletal and clayey; lithic
142. Haplustolls-Calciustolls-Chromusterts; undulating-hilly; clayey-skeletal and clayey; lithic
143. Haplustolls-Ustochrepts-Haplustalfs; undulating-hilly; clayey-skeletal, loamy, and clayey; lithic
144. Paleustalfs-Haplustalfs; undulating-rolling; clayey and fine-loamy
145. Paleustalfs-Rhodustalfs, shallow-Haplustalfs; undulating-hilly; clayey and loamy
146. Paleustalfs-Ustochrepts, shallow-Argialbolls; level-rolling; clayey
147. Paleustalfs-Ustochrepts-Ustipsamments; undulating-rolling; loamy and sandy
148. Paleustolls-Natrustolls-Ustochrepts; level-rolling; clayey
149. Paleustolls-Paleustalfs; level-undulating; clayey, fine-loamy and coarse-loamy
150. Ustochrepts; undulating-hilly; loamy and clayey; shallow
151. Ustochrepts-Argiustolls-Haplustalfs; level-rolling; loamy and clayey; shallow
152. Ustochrepts-Argiustolls-Paleustolls; level-hilly; clayey and loamy; shallow
153. Ustochrepts-Argiustolls-Ustifluvents; level; loamy and sandy

ARIDIC USTOLLS AND USTALFS, THERMIC

154. Calciustolls-Argiustolls-Ustochrepts; level-rolling; loamy and clayey
155. Calciustolls-Paleorthids-Rock land; undulating-steep; loamy-skeletal and loamy; lithic
156. Calciustolls-Paleustolls; level-undulating; fine-loamy and clayey; shallow
157. Calciustolls-Ustochrepts-Calciorthids, shallow, loamy; undulating-hilly; fine-loamy
158. Paleustalfs-Haplustalfs-Ustochrepts; level-rolling; fine-loamy, coarse-loamy, and clayey
159. Paleustalfs-Paleustolls-Calciustolls; level-undulating; fine-loamy
160. Paleustalfs-Ustipsamments; level-undulating; fine-loamy, coarse-loamy, and sandy
161. Paleustolls-Calciustolls; level-undulating; clayey and fine-loamy

USTIC ARIDISOLS AND TORRIORTHENTS, THERMIC

162. Calciorthids-Haplargids-Paleustalfs; undulating-rolling; loamy and loamy-skeletal; shallow
163. Calciorthids-Paleorthids-Calciustolls; undulating-rolling; loamy and loamy-skeletal; lithic
164. Calciorthids-Paleorthids-Camborthids; level-rolling; loamy and clayey; shallow
165. Haplargids-Calciorthids-Torrerts; level-rolling; clayey and fine-silty
166. Haplargids-Camborthids-Torriorthents; level-undulating; clayey, fine-loamy, and fine-silty
167. Haplargids-Torripsamments; level-undulating; fine-loamy and sandy
168. Torriorthents-Camborthids-Haplargids; undulating-hilly; loamy, loamy-skeletal, and clayey; lithic

TYPIC ARIDISOLS AND TORRIORTHENTS, THERMIC

169. Gypsiorthids-Calciorthids-Paleorthids; level-undulating; clayey and loamy
170. Torriorthents-Paleorthids-Rock land; undulating-hilly; loamy-skeletal and loamy; lithic

TYPIC USTOLLS, USTALFS, AND AQUALFS, HYPERTHERMIC

171. Calciustolls-Haplustolls-Ustochrepts; level-undulating; clayey and coarse-loamy
172. Haplustolls-Argiustolls-Haplustalfs; level-undulating; fine-loamy
173. Ochraqualfs-Paleustolls-Argiustolls; level-undulating; clayey and fine-loamy
174. Paleustalfs-Haplustalfs; level-undulating; clayey, sandy and fine-loamy
175. Paleustalfs, arenic, loamy-Ustipsamments; level-undulating; sandy
176. Paleustolls-Pellusterts-Paleorthids, shallow, loamy; level-undulating; clayey

ARIDIC USTOLLS, USTALFS, AND USTOCHREPTS, HYPERTHERMIC

177. Haplustalfs-Paleorthids, shallow, loamy; level-undulating; fine-loamy and clayey
178. Haplustolls-Pellusterts-Paleorthids, shallow, loamy; level-undulating; clayey
179. Paleustalfs-Paleorthids, shallow, loamy; level-undulating; fine-loamy
180. Ustochrepts-Haplustalfs; level-undulating; fine-loamy

PSAMMENTS

181. Psamments; undulating-rolling; sandy
182. Torripsamments-Haplargids-Paleargids; level-rolling; sandy, coarse-loamy, and fine-loamy
183. Ustipsamments-Paleargids; undulating-rolling; sandy, coarse-loamy, and fine-loamy

ROCK LAND

184. Badlands-Torriorthents; undulating-steep; loamy and clayey
185. Rock land; undulating-steep
186. Rock land-Haplargids-Torriorthents; undulating-steep; loamy; lithic
187. Rock land-Torriorthents; undulating-steep; sandy-skeletal and loamy

VERTISOLS

188. Chromusterts-Haplustolls-Ustochrepts; level; clayey and loamy; fluventic; hyperthermic
189. Pellusterts-Chromusterts-Calciustolls; level-unduiating; clayey and fine-silty; thermic
190. Pellusterts-Chromusterts-Paleustalfs; level-undulating; clayey; thermic
191. Pellusterts-Ochraqualfs-Argiustolls; level; clayey and fine-loamy; hyperthermic
192. Pellusterts-Paleorthids, shallow, loamy; level-undulating; clayey; hyperthermic
193. Pellusterts-Paleustolls-Haplustolls; level-undulating; clayey; thermic
194. Pellusterts- Udipsamments-Salorthids; level-undulating; clayey, loamy, and sandy; hyperthermic
195. Torrerts-Pellusterts-Paleorthids, shallow, loamy-skeletal; level-undulating; clayey; hyperthermic

EXPLANATION OF PATTERNS AND SPECIAL SYMBOLS

- Dominantly wet soils.
- More than 20 percent shallow or lithic soils, hilly or steeper soils, but not dominantly Rock land.
- Dominantly Alfisols.
- More than 20 percent Vertisols or Vertic soils but not dominantly Vertisols.

The gray pattern, in the above blocks, is used to represent the color patterns, on the map, which indicate differences of soil temperature and soil moisture.

In areas with more than one of the above characteristics, only the most important is indicated.

- About 100 square miles of Natriborolls, Natrustolls, or Natrargids.
- Large hill or small mountain.
- Narrow areas of level soils, 1 to 3 miles wide, are common along streams and rivers.

CANADA

MEXICO

GULF OF MEXICO

USE OF SOIL MAPS

WISE DECISIONS OF THE USE OF LAND REQUIRE FULL USE OF SOILS INFORMATION. THIS MAP IS USEFUL FOR GENERAL PLANNING AND EDUCATION. MORE SPECIFIC DECISIONS REQUIRE THE USE OF MAPS WHICH SHOW MORE DETAIL SUCH AS STATE OR COUNTY MAPS. SOME DECISIONS, SUCH AS LOCATION AND CONSTRUCTION OF BUILDINGS, DAMS, AND HIGHWAYS, REQUIRE ON-SITE INVESTIGATIONS.

EXPLANATION OF THE LEGEND

The legend is based upon the soil classification system adopted by the National Cooperative Soil Survey, January 1965. Soil temperature and soil moisture are important characteristics used in the system and they are given major emphasis in the legend. The colors on the map, except pink, red, and purple, show differences in soil temperature or soil moisture. Areas on the map dominated by Psamments - sandy soils - are shown in pink; areas dominated by Vertisols - clayey soils are purple; and areas dominated by rocks are red.

The names used in the soil classification system are explained below the map.

Yellow indicates areas dominated by Typic Aridisols, usually dry soils low in organic matter but with pedogenic horizons. These horizons or layers resulted from the changes which occurred as the soils formed. Typic Aridisols are the driest Aridisols and in the Great Plains all of them have thermic temperatures.

Blue indicates areas dominated by Borolls which have mean annual soil temperatures less than 47°F (8°C), frigid. Dark blue indicates Udic Borolls which are more moist than average for Borolls. Medium blue indicates Typic Borolls with average moisture. Light blue indicates Aridic Borolls which are less moist than average. Bluish yellow indicates areas dominated by Borollic Aridisols which also have frigid temperatures but are lighter colored and lower in organic matter than Borolls.

Green indicates areas dominated by Ustolls or Udolls with mesic soil temperatures, those with mean annual soil temperatures between 47°F (8°C) and 59°F (15°C). Dark green indicates Udolls which are usually moist. Ustolls are less moist than Udolls. Moderately dark green indicates Udic Ustolls which are more moist than average for Ustolls. Moderately light green indicates Typic Ustolls with average moisture. Light green indicates Aridic Ustolls which are less moist than average. Greenish yellow indicates areas dominated by Ustic Aridisols with mesic soil temperatures.

Reddish brown indicates areas dominated by Ustolls, Ustalfs, or Udolls with thermic soil temperatures, those with mean annual soil temperatures between 59°F (15°C) and 72°F (22°C). Dark reddish brown indicates Udolls. Moderately dark reddish brown indicates Udic Ustolls and Ustalfs which are more moist than average. Moderately light reddish brown indicates Typic Ustolls and Ustalfs with average moisture. Light reddish brown indicates Aridic Ustolls and Ustalfs which are less moist than average. Light brownish yellow indicates areas dominated by Ustic Aridisols with thermic temperatures.

Orange indicates areas dominated by Ustolls and Ustalfs with hyperthermic soil temperatures, those with mean annual soil temperatures of 72°F (22°C) or higher. Dark orange indicates Typic Ustolls and Ustalfs and light orange indicates Aridic Ustolls and Ustalfs.

Mapping units (areas shown on the soil map) are further defined on the bases of great groups, slope gradients, and particle-size classes. Other properties such as depth to bedrock, excess salt, and active clays also are used where important. The great groups used are briefly described below the map where the names used in the soil classification system are explained.

If one great group includes more than 70 percent of an area, only that one is used. If two great groups include more than 70 percent, only two are used. If three or more great groups are needed to include more than 70 percent, only three great groups are used. A miscellaneous land type - Rock land - is used with great groups of mapping units which include portions dominated by rocks. Badlands with areas of barren soft shales are an example.

The slope range which includes more than 70 percent of the area of a mapping unit is given using one or more of the following terms:

Level	0-3 percent slope
Undulating	3-8 percent slope
Rolling	8-16 percent slope
Hilly	16-30 percent slope
Steep	more than 30 percent slope

As an example "level-undulating" means that more than 70 percent of the area has slope gradients of 0-8 percent. "Undulating-rolling" means that the slope gradients of more than 70 percent of the area are between 3 and 16 percent.

The particle-size classes are listed in the order of their extent until their combined acreage exceeds 70 percent of the mapping unit.

The particle-size classes are those used in the definition of soil families and they apply to the whole soil which includes both the rock fragments and the fine earth. Rock fragments include those mineral particles and loose rocks (not bedrocks) larger than 2 millimeters (1 millimeter equals .04 inch), and the fine earth includes those smaller than 2 millimeters (mm). Soil texture classes apply only to the fine earth and they are not used in this legend.

The particle-size class generally is that of the 10 to 40-inch layer. The particle-size classes used in this legend are as follows:

Sandy	Fine earth includes mainly fine sand (0.25 - 0.1 mm) and coarser, some very fine sand (0.1 - 0.05 mm) but very little silt (0.05 - 0.002 mm) and clay (less than 0.002 mm). Rock fragments are less than 35 percent by volume.
Loamy	Fine earth includes less than 35 percent clay, variable amounts of silt and sands and usually less than 70 percent fine sands and coarser. Rock fragments are less than 35 percent by volume.
Coarse-loamy	A loamy particle-size with less than 18 percent clay and more than 15 percent fine sand and coarser in the fine earth.
Fine-loamy	A loamy particle-size with 18-35 percent clay and more than 15 percent fine sand and coarser in the fine earth.
Coarse-silty	A loamy particle-size with less than 18 percent clay and less than 15 percent fine sand and coarser in the fine earth.
Fine-silty	A loamy particle-size with 18-35 percent clay and less than 15 percent fine sand and coarser in the fine earth.
Clayey	Fine earth contains more than 35 percent clay. Rock fragments are less than 35 percent by volume.
Sandy-skeletal	Rock fragments are 35 percent or more by volume and fine earth is sandy.
Loamy-skeletal	Rock fragments are 35 percent or more by volume and fine earth is loamy.
Clayey-skeletal	Rock fragments are 35 percent or more by volume and fine earth is clayey.
Fine-loamy over sandy or sandy-skeletal	Contrasting particle-sizes with the upper part of the 10 to 40-inch layer fine-loamy and the lower part sandy or sandy-skeletal.

Additional properties used in some mapping units are as follows:

Lithic	Soils with hard bedrock within 20 inches of the surface occupy more than 20 percent of the area.
Shallow	Soils with soft bedrock within 20 inches of the surface occupy more than 20 percent of the area.
Saline	Soils with excess salts dominate these areas.
Vertic	Soils with very active or swelling clays occupy more than 20 percent of the area.
Fluventic	Soils of the flood plains formed in alluvium occupy more than 20 percent of the area.
Arenic	Soils with a 20 to 40-inch surface horizon that is sandy throughout.

If any of the above properties apply to only one great group, they are shown immediately after that great group. For example, in mapping unit 122 only the Torriorthents are lithic and loamy.

ACKNOWLEDGEMENT

The map was prepared by Andrew R. Aandahl, Professor of Agronomy, UNIVERSITY OF NEBRASKA, and for many years, Principal Soil Correlator for the Great Plains States and later the Midwest Region, SOIL CONSERVATION SERVICE, U. S. DEPARTMENT OF AGRICULTURE. The map was prepared in cooperation with soil specialists in the Great Plains and with the help of an advisory committee of state and federal soil specialists.

STATE AGRICULTURAL EXPERIMENT STATIONS, COOPERATIVE EXTENSION SERVICES, the AGRICULTURAL RESEARCH SERVICE, and the SOIL CONSERVATION SERVICE contributed to the project. The GREAT PLAINS AGRICULTURAL COUNCIL provided valuable assistance.

The cartographic work was provided by the SOIL CONSERVATION SERVICE.

The data for the map were from published and unpublished soil maps and reports.

The map is part of a project which also includes preparation of a book on soil-agriculture relationships in the Plains.

© Copyright Andrew R. Aandahl 1972

Copies of the map may be obtained from:

Andrew R. Aandahl
P.O. Box 81242
Lincoln, Neb. 68508

SCALE: 1/2,500,000

SCALE 25 0 25 50 75 100 MILES

ALBERS EQUAL-AREA PROJECTION
NORTH AMERICAN DATUM

LOCATION MAP

GREAT PLAINS

country 108
subject soils
scale 1:2,500,000
map rel. 3
libr. rel. 225

EXPLANATION OF THE NAMES

The names used in the soil classification system are combinations of formative elements from Latin and Greek. If the formative elements had been taken from English, confusion would have resulted because English words commonly have more than one meaning. The formative elements have specific meaning in the system as shown in the table.

The soil classification system consists of orders which are subdivided into suborders, which are subdivided into great groups, which are subdivided into subgroups, which are subdivided into families which are subdivided into series. Only names of suborders and great groups are used in the legend.

The names of orders end in "sol" and each order name contains a formative element which is the final syllable of the names of suborders, great groups, and subgroups of the order. The orders with their formative element and a brief description are as follows:

Alfisols	alf	Soils with gray to brown surface horizons, subsurface horizons of clay accumulation, medium to high in bases.
Aridisols	id	Soils that are low in organic matter, usually dry, and have pedogenic horizons.
Entisols	ent	Soils without pedogenic horizons.
Histosols	ist	Organic soils formed from plant residues.
Inceptisols	ept	Soils with faint or indistinct pedogenic horizons and are usually moist.
Mollisols	oll	Soils with nearly black surface horizons high in organic matter and bases.
Oxisols	ox	Soils that are mixtures principally of kaolin, hydrated oxides, and quartz.
Spodosols	od	Soils that have amorphous materials consisting of organic matter plus compounds of aluminum and usually iron accumulated in a subsurface horizon.
Ultisols	ult	Soils that have subsurface horizons of clay accumulation and are low in bases.
Vertisols	ert	Soils that are clayey and have wide deep cracks when dry.

Pedogenic horizons result from the changes that occur in the soils as they form. The more common bases are calcium, magnesium, potassium, and sodium.

Histosols, Oxisols, Spodosols, and Ultisols are not in this legend.

The names of suborders consist of two syllables, the formative element of the order plus another formative element used as a prefix. Aquolls, for example, are wet Mollisols.

The names of great groups consist of the suborder names with another formative element added as a prefix. The letter "o" or "i" is sometimes added between the formative element and the suborder name to make the great group name easier to pronounce.

The formative elements used in this legend to form suborder and great group names and their meaning are as follows:

Alb	Soils that have a bleached (white) surface or subsurface horizon reflecting wetness.
Aqu	Soils that are wet for long periods unless artificially drained.
Arg	Soils with a horizon of clay accumulation.
Bor	Soils with mean annual temperatures less than 47°F (8°C).
Calc	Soils that have a horizon of calcium carbonate accumulation and are calcareous throughout.
Camb	Soils with minor or slight changes or alterations.
Chrom	Soils with a brownish or reddish surface horizon.
Eutro	Fertile soils high in bases.
Fluv	Soils in flood plains formed in alluvium.
Gyps	Soils that have a horizon of gypsum, calcium sulfate.
Hapl	The typical ones used in the names of great groups.
Natr	Soils with an accumulation of clay high in sodium.
Ochr	Soils with a light-colored surface horizon low in organic matter.
Orth	The typical ones used in the names of suborders.
Pale	Soils that are old and on stable landscapes, which have not been disturbed by recent geologic erosion or depositions.
Pell	Soils that have a black or dark gray surface horizon.
Psamm	Soils that are sandy with very little silt and clay.
Rhod	Soils with a dark red color.
Sal	Soils with an accumulation of easily soluble salts.
Torr	Soils that are usually dry.
Ud	Soils that are usually moist and have long periods with enough moisture for plant growth.
Ust	Soils that are sometimes dry, sometimes moist, and have summer rains but lack moisture for plant growth for long periods.

The legend includes 16 suborders and 41 great groups.

The suborders of alfisols are as follows:

- Aqualfs — Alfisols that are wet for long periods.
- Boralfs — Alfisols with mean annual soil temperatures less than 47°F (8°C).
- Ustalfs — Alfisols that are sometimes dry, sometimes moist, have summer rains, lack moisture for plant growth for long periods and have mean annual soil temperatures of 47°F (8°C) or higher.

Albaqualfs and Ochraqualfs are great groups of Aqualfs. Albaqualfs have a more abrupt increase in clay content between the light colored horizon and the horizon of clay accumulation than Ochraqualfs.

Eutroboralfs are the only great group of Boralfs represented and they are fertile soils high in bases.

The Ustalfs are represented by four great groups; namely,

- Haplustalfs which are the typical Ustalfs which are free or nearly free of sodium, not red or old.
- Natrustalfs which have much sodium on the clays.
- Rhodustalfs which have thin dark red horizons of clay accumulation.
- Paleustalfs which are old soils with thick red horizons of clay accumulation, and occur on old stable landscapes.

The suborders of Aridosols are as follows:

- Orthids - The Aridisols without a horizon of clay accumulation.
- Argids - The Aridisols with a horizon of clay accumulations.

The Orthids are represented by five great groups; namely,

- Calciorthids which are calcareous throughout and have a distinct horizon of calcium carbonate.
- Camborthids which have had only slight changes during their formation and no large accumulations of calcium carbonates or soluble salts.
- Gypsiorthids which have a distinct horizon of calcium sulfate, gypsum.
- Paleorthids which are old soils with a very hard, usually thick horizon cemented by calcium carbonates.
- Salorthids which have large amounts of easily soluble salts and the ground water is high enough for long enough period to maintain their salt content.

The Argids are represented by three great groups; namely,

- Haplargids which have little or no sodium on the clays and are not old, the typical ones.
- Natrargids which have much sodium on the clays.
- Paleargids which are old soils with either a very hard horizon cemented by calcium carbonates or an abrupt increase in clay content in the upper 3 inches of the horizon of clay accumulation.

The suborders of Entisols are as follows:

- Fluvents which are soils of the flood plain formed in alluvium.
- Orthents which are not sandy or in the flood plain, the typical ones.
- Psamments which are sandy.

The Fluvents are represented by two great groups; namely,

- Ustifluvents which are sometimes moist, sometimes dry.
- Torrifluvents which are usually dry.

The Orthents are represented by three great groups; namely,

- Udorthents which are of humid climates.
- Ustorthents which are sometimes moist, sometimes dry.
- Torriorthents which are of dry climates.

The Psamments are represented by three great groups; namely,

- Udipsamments which are of humid climates.
- Ustipsamments which are sometimes moist, sometimes dry.
- Torripsamments which are of dry climates.

The only suborder of Inceptisol used in the legend is Ochrepts which have a light-colored, low-organic matter surface horizon. Ustochrepts is the only great group of Ochrepts and these are sometimes dry, sometimes moist.

The suborders of Mollisols are as follows:

- Albolls which have a mottled, bleached, or whitish subsurface horizon over a mottled or gray horizon of clay accumulation. Mottles are small spots of contrasting colors usually evidence of wetness.
- Aquolls which are wet, usually for long periods.
- Borolls which have mean annual soil temperatures less than 47°F (8°C).
- Udolls which are usually moist and have long periods with enough moisture for plant growth and have a mean annual soil temperature of 47°F (8°C) or higher.
- Ustolls which are sometimes dry, sometimes moist, have long periods without enough water for plant growth, summer rains, mean annual soil temperatures of 47°F (8°C) or higher and usually some accumulation of calcium carbonate in the lower part of the profile.

Albolls are represented by only one great group, Argialbolls which have a clay accumulation nearly free of sodium.

The Aquolls are represented by three great groups; namely,

- Argiaquolls which have a horizon of clay accumulation.
- Calciaquolls which are calcareous throughout and have a large accumulation of calcium carbonate near the surface.
- Haplaquolls which lack a large accumulation of clay or calcium carbonate, the typical ones.

The Borolls are represented by four great groups; namely,

- Argiborolls which have a horizon of clay accumulation.
- Calciborolls which are calcareous throughout and have large accumulations of calcium carbonate.
- Haploborolls which lack a distinct accumulation of calcium carbonate, clay or sodium, the typical ones.
- Natriborolls which have a horizon of clay accumulation with much sodium on the clays.

The Udolls are represented by three great groups; namely,

- Argiudolls which have a horizon of clay accumulation.
- Hapludolls which are the typical ones and lack a horizon of clay accumulation or age.
- Paleudolls which are old soils with thick reddish horizons of clay accumulation.

The Ustolls are represented by five great groups; namely,

- Argiustolls which have a horizon of clay accumulation.
- Calciustolls which are calcareous throughout and have a large accumulation of calcium carbonate.
- Haplustolls which lack a distinct accumulation of calcium carbonate, clay or sodium, the typical ones.
- Natrustolls which have a horizon of clay accumulation with much sodium on the clays.
- Paleustolls which are old soils of stable landscapes with thick reddish horizons of clay accumulation, a very hard horizon cemented by calcium carbonate or an abrupt increase in clay content in the upper 3 inches of the horizon of clay accumulation.

The suborder of Vertisols are as follows:

- Usterts which are sometimes dry, sometimes moist, and have summer rains.
- Torrerts which are usually dry.

The Usterts are represented by two great groups; namely,

- Pellusterts which are black or dark gray.
- Chromusterts which are brownish.

All Torrerts are so similar that they are not divided into great groups.